

Question 1

Q1	Model Solution – 30 Marks	Marking Notes
(a)	$\frac{(4-2i)}{(2+4i)} = \frac{4-2i}{2+4i} \times \frac{2-4i}{2-4i}$ $= \frac{(8-16i-4i-8)}{2^2+4^2}$ $= \frac{-20i}{20}$ $= 0 - 1i$ $\therefore k = -1$ <p style="text-align: center;">OR</p> $\frac{(4-2i)}{(2+4i)} = \frac{-i(2+4i)}{2+4i}$ $= 0 - 1i$ $\therefore k = -1$ <p style="text-align: center;">OR</p> $4 - 2i = ki(2 + 4i)$ <p>Re: $4 = -4k \quad \therefore k = -1$</p> <p>or Im: $-2i = 2ki \quad \therefore k = -1$</p>	<p>Scale 10C(0, 3, 7, 10)</p> <p><i>Note:</i> Accept $0 - 1i$</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • $2 - 4i$ • $4 - 2i = ki(2 + 4i)$ <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • $\frac{4-2i}{2+4i} \times \frac{2-4i}{2-4i}$ • $4 - 2i = -4k + 2ki$ • Sets $\text{Re} = \text{Re}$ or $\text{Im} = \text{Im}$ <p><i>Full Credit –1:</i></p> <ul style="list-style-type: none"> • $0 - i$ or $-1i$ as solution, with k not identified.
Q1	Model Solution – 30 Marks	Marking Notes
(b)	$-5 + 12i = (a + bi)^2$ $a^2 + 2abi - b^2$ <p>Re: $a^2 - b^2 = -5$</p> <p>Im: $2ab = 12 \quad \therefore b = \frac{6}{a}$</p> $a^2 - \left(\frac{6}{a}\right)^2 = -5$ $a^4 + 5a^2 - 36 = 0$ $(a^2 + 9)(a^2 - 4) = 0$ <p>$\therefore a = \pm 2$ and $b = \pm 3$</p> <p>Answer: $2 + 3i, -2 - 3i$</p> <p style="text-align: center;">OR</p> $r = \sqrt{5^2 + 12^2} = 13$ $\tan \theta = -\frac{12}{5} \text{ so } \cos \theta = -\frac{5}{13}$ $(-5 + 12i)^{\frac{1}{2}}$ $= [13(\cos(\theta + 2n\pi) + i \sin(\theta + 2n\pi))]^{\frac{1}{2}}$ $= \sqrt{13} \left(\cos\left(\frac{\theta}{2} + n\pi\right) + i \sin\left(\frac{\theta}{2} + n\pi\right) \right)$ $2 \sin^2\left(\frac{\theta}{2}\right) = 1 - \cos \theta = 1 + \frac{5}{13}$	<p>Scale 10D(0, 3, 5, 8, 10)</p> <p><i>Note:</i> Accept $2 + 3i$ for <i>Full Credit</i></p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • $(a + bi)^2 = -5 + 12i$ • $a + bi = (-5 + 12i)^{\frac{1}{2}}$ • r or θ found • $-5 + 12i$ plotted on Argand diagram. • Shows some knowledge of De Moivre's theorem <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • Relevant equation in a single variable • Writes $-5 + 12i$ in polar form <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • Finds $a = 2$ or $b = 3$ • $-2 - 3i$ found • Correct solution in polar form (accept with mishandling of $2n\pi$)

	<p>So $\sin\left(\frac{\theta}{2}\right) = \frac{3}{\sqrt{13}}$ and so $\cos\left(\frac{\theta}{2}\right) = \frac{2}{\sqrt{13}}$</p> <p>$n = 0$: $\sqrt{13}\left(\cos\left(\frac{\theta}{2}\right) + i\sin\left(\frac{\theta}{2}\right)\right)$ $= \sqrt{13}\left(\frac{2}{\sqrt{13}} + i\frac{3}{\sqrt{13}}\right) = 2 + 3i$</p> <p>$n = 1$: $\sqrt{13}\left(\cos\left(\frac{\theta}{2} + \pi\right) + i\sin\left(\frac{\theta}{2} + \pi\right)\right)$ $= \sqrt{13}\left(-\frac{2}{\sqrt{13}} - i\frac{3}{\sqrt{13}}\right) = -2 - 3i$</p>	
Q1	Model Solution – 30 Marks	Marking Notes
(c)	$z^3 = r(\cos\theta + i\sin\theta)$ $z = (r(\cos\theta + i\sin\theta))^{\frac{1}{3}}$ $= 2\left(\cos\frac{\pi+2n\pi}{3} + i\sin\frac{\pi+2n\pi}{3}\right)$ <p>$n = 0$: $z = 2\left(\cos\frac{\pi}{3} + i\sin\frac{\pi}{3}\right) = 1 + \sqrt{3}i$</p> <p>$n = 1$: $z = 2\left(\cos\frac{3\pi}{3} + i\sin\frac{3\pi}{3}\right) = -2$</p> <p>$n = 2$: $z = 2\left(\cos\frac{5\pi}{3} + i\sin\frac{5\pi}{3}\right) = 1 - \sqrt{3}i$</p>	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p><i>Note:</i> if $((r(\cos\theta + i\sin\theta))^3)$ is used, award <i>Low Partial Credit</i> at most.</p> <p><i>Note:</i> polar form must be used to achieve any credit</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • $z = (r(\cos\theta + i\sin\theta))^{\frac{1}{3}}$ • r found • θ found • $-8 + 0i$ plotted on an Argand diagram • Shows some knowledge of De Moivre's theorem <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • $z = 8^{\frac{1}{3}}\left(\cos\frac{\pi}{3} + i\sin\frac{\pi}{3}\right)$ • $8^{\frac{1}{3}}\left(\cos\frac{\pi+2n\pi}{3} + i\sin\frac{\pi+2n\pi}{3}\right)$ <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • One root evaluated in the form $a + bi$ from De Moivre's expression • Three solutions in polar form


Question 2

Q6	Model Solution – 30 Marks	Marking Notes
(a)	$h'(x) = a(x+1)(x-3)$ $= a(x^2 - 2x - 3)$ $(0, 6) \in h'(x)$ $\therefore 6 = a(0 + 0 - 3)$ $\Rightarrow a = -2$ $h'(x) = -2(x^2 - 2x - 3)$ $h'(x) = -2x^2 + 4x + 6$ <p style="text-align: center;">OR</p> $h'(x) = -2x^2 + 4x + 6$ $\text{y-intercept} = h'(0) = 6$ $h'(x) = -2(x+1)(x-3)$ $\therefore \text{Roots} = -1 \text{ and } 3$ <p style="text-align: center;">OR</p> $h'(x) = ax^2 + bx + c$ $h'(0) = c = 6$ $\text{So } h'(-1) = a - b + 6 = 0$ $\text{and } h'(3) = 9a + 3b + 6 = 0$ $3 \times h'(1) = 3a - 3b + 18 = 0$ $\text{So } 12a + 24 = 0$ $\therefore a = -2 \text{ and } b = 4$ $\text{i.e. } h'(x) = -2x^2 + 4x + 6$	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p><i>Note:</i> Accept 3 points from graph verified as belonging to given equation of $h'(x)$</p> <p><i>Note:</i> three points identified from graph is <i>Low Partial Credit</i>.</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> Identifies or uses a relevant value from the graph, for example, $(x + 1)$ or 3 A factorisation of given $h'(x)$, for example $2(-x^2 + 2x + 3)$ <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> Generates $(x + 1)(x - 3)$ from graph Correctly verifies one point from the graph into the given $h'(x)$ Full factorisation of given $h'(x)$ Using simultaneous equations, finds one value (a, b, or c) <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> Generates $x^2 - 2x - 3$ from graph From given $h'(x)$, finds two roots Correctly verifies two points from the graph into the given $h'(x)$ From given $h'(x)$, shows that y-intercept is 6 and fully factorises Using simultaneous equations, finds two values (from a, b, and c)
Q6	Model Solution – 30 Marks	Marking Notes
(b)	$h''(x) = -4x + 4 = 0 \text{ at max/min of } h'(x)$ $\therefore x = 1$ $h'''(x) = -4 < 0, \text{ i.e. max}$ $h'(1) = -2(1)^2 + 4(1) + 6 = 8$ <p style="text-align: center;">OR</p> <p>[Quadratic with negative x^2, so max occurs halfway between the roots:]</p> $x = \frac{-1+3}{2} = 1$ $h'(1) = -2(1)^2 + 4(1) + 6 = 8$ <p style="text-align: center;">OR</p> $h'(x) = -2(x^2 - 2x - 3)$ $= -2(x^2 - 2x + 1 - 1 - 3)$ $= -2((x - 1)^2 - 4)$ $= -2(x - 1)^2 + 8$	<p>Scale 10C(0, 3, 7, 10)</p> <p><i>Note:</i> It is possible to accept for <i>Full Credit</i> without $h'''(x) < 0$</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> Some correct differentiation of $h'(x)$ Finds $h''(x)$ $h'(x) = -2(x^2 - 2x - 3)$ Indicates axis of symmetry on graph <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> $x = 1$ $h'(x) = -2((x - 1)^2 - 4)$

	∴ max positive slope= 8	
(c)	$h(x) = \int h'(x) dx$ $h(x) = -\frac{2x^3}{3} + \frac{4x^2}{2} + 6x + C$ <p style="text-align: center;">(0, -2) ∈ h(x):</p> $-2 = -0 + 0 + 0 + C$ $\Rightarrow C = -2$ $\therefore h(x) = -\frac{2x^3}{3} + 2x^2 + 6x - 2$	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p><i>Note:</i> Accept correct answer without work.</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Any indication of integration <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • Integration of 3 terms fully correct <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • Relevant equation in C (with substitution)

Question 3

Q2	Model Solution – 30 Marks	Marking Notes
(a)	$f'(x) = 2x + b$ $f'(3) = 2(3) + b = 0$ $b = -6$ $f(3) = (3)^2 - 6(3) + c = -1$ $9 - 18 + c = -1$ $c = 8$ <p style="text-align: center;">OR</p> $x^2 + bx + c = \left(x + \frac{b}{2}\right)^2 - \frac{b^2}{4} + c$ $-\frac{b}{2} = 3 \text{ so } b = -6$ $-\frac{b^2}{4} + c = -1 \text{ so } c = 8$ <p style="text-align: center;">OR</p> $f(x) = (x - 3)^2 - 1$ $= x^2 - 6x + 8$	<p>Scale 15D (0, 4, 8, 12, 15)</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit, for example, $f(3)$ or some correct differentiation • Work of merit at completing the square • $(x - h)^2 + k$ <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • b correct • Uses $f(3)$ to find a correct equation in b and c • $\left(x + \frac{b}{2}\right)^2 - \frac{b^2}{4} + c$ • Work of merit in finding both b and c • $(x - 3)^2 + k$, where $k \neq -1$ • $(x - h)^2 - 1$, where $h \neq 3$ <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> • Finds b and work of merit in finding c • $(x - 3)^2 - 1$
(b)	$\lim_{n \rightarrow \infty} \left(\frac{n}{n+1} + \frac{n+1000}{n} + \left(\frac{1}{3}\right)^n \right)$ $= \lim_{n \rightarrow \infty} \left(\frac{n}{n+1} \right) + \lim_{n \rightarrow \infty} \left(\frac{n+1000}{n} \right) + \lim_{n \rightarrow \infty} \left(\left(\frac{1}{3}\right)^n \right)$ $\left[= \lim_{n \rightarrow \infty} \left(\frac{1}{1 + \frac{1}{n}} \right) + \lim_{n \rightarrow \infty} \left(\frac{1 + \frac{1000}{n}}{1} \right) + \lim_{n \rightarrow \infty} \left(\left(\frac{1}{3}\right)^n \right) \right]$ $= \frac{1}{1+0} + \frac{1+0}{1} + 0$ $= 2$	<p>Scale 5C (0, 2, 3, 5)</p> <p>Note: Full credit for correct answer without work.</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit, for example, indicates sum of limits, divides by highest power of n in one of first two terms • Substitutes ∞ for n • Finds two or more terms of the sequence, $T_n = \frac{n}{n+1} + \frac{n+1000}{n} + \left(\frac{1}{3}\right)^n$ <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • One limit correctly evaluated and work of merit in any one of the other two limits

Q2	Model Solution – 30 Marks	Marking Notes
(c)	 <p>The figure consists of two coordinate planes, each with x and y axes ranging from -5 to 5. The top graph shows a grey curve labeled $g(x)$ with a vertex at $(-1, 1)$. A black curve is drawn, which is a vertical stretch and reflection of $g(x)$, with its vertex at $(-1, -1)$. The bottom graph shows the same grey curve $g(x)$. A black curve is drawn, which is a horizontal stretch and reflection of $g(x)$, with its vertex at $(-4, 1)$.</p>	<p>Scale 10C (0, 4, 7, 10)</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit in one part, for example, one point correctly transformed • In part (i) any vertical translation of $g(x)$ • In part (ii) any horizontal translation of $g(x)$ • In part (i) finds $g(x + 2)$ or $g(x - 2)$ • In part (ii) finds $g(x) - 3$ or $g(x) + 3$ <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • One part correct • Work of merit in both parts

Question 4

Q3	Model Solution – 30 Marks	Marking Notes
(a)	<p>Assume that $\sqrt{2}$ is rational. $\sqrt{2} = \frac{a}{b}$ where $a, b \in \mathbb{Z}, b \neq 0$ and $\text{HCF}(a, b) = 1$ $2 = \frac{a^2}{b^2}$ $2b^2 = a^2$ $\Rightarrow a^2$ is even If a^2 is even, then a is even. $\therefore a = 2k$, where $k \in \mathbb{Z}$ $2b^2 = (2k)^2$ $2b^2 = 4k^2$ $b^2 = 2k^2$ $\therefore b^2$ is even If b^2 is even, then b is even. If both a and b are even, then they have 2 as a common factor. This contradicts the assumption that $\text{HCF}(a, b) = 1$.</p>	<p>Scale 10D (0, 3, 5, 8, 10) <i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit, for example $\sqrt{2} = \frac{a}{b}$ • Work of merit in showing that a is even <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • Shows that a is even <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> • Shows both a and b are even
(b)	<p>Method 1</p> $\log_3 t + \frac{\log_3 t}{\log_3 9} + \frac{\log_3 t}{\log_3 27} + \frac{\log_3 t}{\log_3 81} = 10$ $\log_3 t + \frac{\log_3 t}{2} + \frac{\log_3 t}{3} + \frac{\log_3 t}{4} = 10$ $12\log_3 t + 6\log_3 t + 4\log_3 t + 3\log_3 t = 120$ $25\log_3 t = 120$ $\log_3 t = \frac{120}{25}$ $t = 3^{\frac{120}{25}} = 3^{\frac{24}{5}}$ <p style="text-align: center;">OR</p> <p>Method 2</p> $\frac{1}{\log_t 3} + \frac{1}{\log_t 9} + \frac{1}{\log_t 27} + \frac{1}{\log_t 81} = 10$ $\frac{1}{\log_t 3} + \frac{1}{2\log_t 3} + \frac{1}{3\log_t 3} + \frac{1}{4\log_t 3} = 10$ $\frac{25}{12\log_t 3} = 10$ $\log_t 3 = \frac{25}{120}$ $t^{\frac{25}{120}} = 3$ $t = 3^{\frac{120}{25}}$	<p>Scale 10D (0, 3, 5, 8, 10) 3 steps: 1. Changing all to the same base 2. Simplifies to an equation in t with one log 3. Finds t</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit, for example, changes the base of one log (from the given equation) • Writes either 9, 27 or 81 in the form 3^k <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • One correct step <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> • 2 correct steps

Q3	Model Solution – 30 Marks	Marking Notes
(c) (i) (ii)	<p>(i) Any valid explanation, for example: the power you need to raise 6 to, to get m.</p> <p>(ii) $\log_6 m > 1$</p>	<p>Scale 10C (0, 4, 7, 10)</p> <p>Note: Accept $6^x = m$ as a valid explanation for (i)</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit in (i) or (ii), for example, some reference to indices • $\log_6 m > 0$ or $\log_6 m$ is positive <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> • (i) or (ii) correct

Question 5

Q10	Model Solution – 50 Marks	Marking Notes
(a)		<p>Scale 5B (0, 2, 5)</p> <p><i>Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit, for example, each rectangle of height 2 units
(b)	<p>Method 1</p> $h = \frac{8}{3}$ <p>Using similar triangles</p> $\frac{w_1}{2} = \frac{8}{3} \div 8 \quad (w_1 = \text{length of middle rectangle})$ $w_1 = \frac{2}{3}$ $w_2 = \frac{4}{3} \quad (w_2 = \text{length of top rectangle})$ $\text{Area} = 2 \left(\frac{8}{3} \right) + \frac{4}{3} \left(\frac{8}{3} \right) + \frac{2}{3} \left(\frac{8}{3} \right)$ $= \frac{32}{3} \text{ units}^2$ <p>Method 2</p> <p>6 small triangles of length $\frac{2}{6} = \frac{1}{3}$</p> $\text{Area}_{\text{small } \Delta's} = 6 \times \frac{1}{2} \times \frac{1}{3} \times \frac{8}{3} = \frac{8}{3}$ $\text{Area}_{\text{big } \Delta} = \frac{1}{2} \times 2 \times 8 = 8$ $\text{Area}_{\text{rectangles}} = \frac{8}{3} + 8 = \frac{32}{3} \text{ units}^2$ <p>Method 3</p> <p>Sum of lengths of horizontal sides of small triangles (i.e., excess of rectangles over large triangle) = 2</p> <p>Height of each small triangle = $\frac{8}{3}$</p> $\sum \text{areas of small } \Delta's = \frac{1}{2} \times 2 \times \frac{8}{3} = \frac{8}{3}$ $\text{Area of large triangle} = \frac{1}{2} \times 2 \times 8 = 8$ $\therefore \text{Area of rectangles} = \frac{8}{3} + 8 = \frac{32}{3}$	<p>Scale 10C (0, 4, 7, 10)</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit in finding dimensions or area of one rectangle or one small triangle, • Finds $h = \frac{8}{3}$ • Finds the area of the large triangle <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • w_1 and w_2 found • Areas of 2 rectangles found • Sum of the areas of the small triangles found with work shown

Q10	Model Solution – 50 Marks	Marking Notes
(c)	<p>Method 1</p> $T_4 = \frac{8}{4} \left[\frac{2}{4} + \frac{4}{4} + \frac{6}{4} + \frac{8}{4} \right]$ <p>Similarly,</p> $T_n = \frac{8}{n} \left[\frac{2}{n} + \frac{4}{n} + \dots + \frac{2n}{n} \right]$ $= \frac{8}{n^2} [2 + 4 + 6 + \dots + 2n]$ <p>$2 + 4 + \dots + 2n$ is an A.P. with $a = 2$ and $d = 2$ and n terms</p> $S_n = \frac{n}{2} [2(2) + (n-1)(2)]$ $= n(n+1)$ $T_n = \frac{8}{n^2} [n(n+1)]$ $= \frac{8}{n} (n+1)$ <p>or equivalent</p> <p>Method 2</p> <p>Total area = area of given triangle plus sum of areas of small triangles.</p> $h = \frac{8}{n}$ <p>$2n$ small triangles</p> <p>Base of each small $\Delta = \frac{2}{2n} = \frac{1}{n}$</p> $\text{Area of small } \Delta's = 2n \times \frac{1}{2} \times \frac{1}{n} \times \frac{8}{n}$ $= \frac{8}{n}$ $\text{Area} = \left(\frac{8}{n}\right) + \left(\frac{1}{2} \times 2 \times 8\right)$ $= \frac{8}{n} + 8$	<p>Scale 5C (0, 2, 3, 5)</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit, for example, work towards establishing pattern by writing $T_k, k \neq 3$ • $\frac{8}{n}$ mentioned as height of rectangle • Identifies $2n$ (or n) small triangles • Base length of each small triangle found <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • $T_n = \frac{8}{n} \left[\frac{2}{n} + \frac{4}{n} + \dots + \frac{2n}{n} \right]$ or equivalent not in closed form • Finds sum of the areas of the small triangles
(d)	<p>$\frac{8(n-1)}{n} > 0.95(8)$</p> $\frac{n-1}{n} > 0.95$ $n-1 > 0.95n$ $0.05n > 1$ $n > 20$ $n = 21$ <p style="text-align: center;">OR</p> $\frac{8n^2(n-1)}{n} > 0.95(8)n^2$ $8n^2 - 7.6n^2 - 8n > 0$ $0.4n^2 - 8n > 0$ $n^2 - 20n > 0$ $n(n-20) > 0$ $n > 20$ $n = 21$	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p>Note: where candidates multiply both sides by n^2, they must find $n = 20$ to be awarded <i>High Partial Credit</i>.</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Work of merit in establishing inequality, for example, finds the area of triangle <p><i>Mid Partial Credit</i></p> <ul style="list-style-type: none"> • Forms the correct inequality <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> • $n - 1 > 0.95n$
(e) (i)	$\int \frac{x^2 c^2}{h^2} dx = \frac{c^2}{h^2} \int x^2 dx$	<p>Scale 10C (0, 4, 7, 10)</p> <p><i>Low Partial Credit:</i></p>

	$= \frac{c^2}{h^2} \left[\frac{x^3}{3} \right]_0^h$ $= \frac{c^2}{h^2} \left[\frac{h^3}{3} - 0 \right]$ $= \frac{c^2 h}{3}$	<ul style="list-style-type: none"> • Integral set up correctly <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> • Integration is correct • Mishandles $\frac{c^2}{h^2}$, but otherwise correct.
Q10	Model Solution – 50 Marks	Marking Notes
(e)(ii)	$\frac{dx}{dt} = 3 \quad \frac{dS}{dt} = \frac{dS}{dx} \times \frac{dx}{dt}$ $S(x) = \frac{x^2 c^2}{h^2}$ $\frac{dS}{dx} = \frac{c^2}{h^2} (2x)$ $\frac{dS}{dt} = \frac{c^2}{h^2} (2x)(3)$ $= \frac{6c^2 x}{h^2}$ <p>When $x = \frac{h}{2}$</p> $\frac{dS}{dt} = \frac{6c^2 \left(\frac{h}{2}\right)}{h^2} = \frac{3c^2}{h}$	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • States a relevant derivative, for example, $\frac{dS}{dx}$ or $\frac{dS}{dt}$ • $x = \frac{h}{2}$ • Some correct differentiation <p><i>Mid Partial Credit</i></p> <ul style="list-style-type: none"> • Any two of the following: <ul style="list-style-type: none"> ○ $\frac{dx}{dt} = 3$ ○ $x = \frac{h}{2}$ ○ $\frac{dS}{dx} = \frac{c^2}{h^2} (2x)$ ○ $\frac{dS}{dt} = \frac{dS}{dx} \times \frac{dx}{dt}$ or similar <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> • $\frac{dS}{dt} = \frac{dS}{dx} \times \frac{dx}{dt}$, and any two others from the MPC list above

Question 6

Q3	Model Solution – 30 Marks	Marking Notes
(a)	$\frac{1}{6} \sin 6x + C$	<p>Scale 5B (0, 2, 5)</p> <p><i>Partial Credit</i></p> <ul style="list-style-type: none"> Some correct integration, for example, $\sin x$ <p><i>Full Credit –1</i></p> <ul style="list-style-type: none"> Apply a * if the +C term is missing
(b) (i)	<p><i>Co-ordinates of the point of contact</i> $f(2) = -21$, so point is $(2, -21)$</p> <p><i>Slope of the tangent at $x = 2$</i> $f'(x) = 6x^2 - 18x + 5$ $f'(2) = -7$... slope of tangent</p> <p><i>Equation of the tangent at $x = 2$</i> $y - (-21) = -7(x - 2)$ or equivalent</p>	<p>Scale 5D (0, 2, 3, 4, 5)</p> <p>Note: Engagement with Step 3 is required to be awarded credit for Step 4</p> <p>Consider solution as involving 4 steps:</p> <ol style="list-style-type: none"> Finds y-value at $x = 2$ Differentiates $2x^3 - 9x^2 + 5x - 11$ Finds $f'(2)$ Finds the equation of the tangent <p><i>Low Partial Credit</i></p> <ul style="list-style-type: none"> Work of merit, for example, some correct differentiation; Substitutes $x = 2$ in $f(x)$; Formula for the equation of a line with some relevant substitution <p><i>Mid Partial Credit</i></p> <ul style="list-style-type: none"> 2 steps correct <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> 3 steps correct
Q3	Model Solution – 30 Marks	Marking Notes
(b) (ii)	$f'(x) = 6x^2 - 18x + 5$ $f''(x) = 12x - 18$ $f''(x) = 0$ $12x - 18 = 0$ $x = \frac{3}{2}$ <p>Also accept the following for Full Credit: <i>x values at local maximum & local minimum</i></p> $f'(x) = 0$ $6x^2 - 18x + 5 = 0$ $x_1 = \frac{18 + \sqrt{204}}{12}$ $x_2 = \frac{18 - \sqrt{204}}{12}$ <p><i>x co-ordinate of the point of inflection:</i></p> $\frac{x_1 + x_2}{2} = \frac{36}{12} \div 2$ 3	<p>Scale 5C (0, 2, 3, 5)</p> <p><i>Low Partial Credit</i></p> <ul style="list-style-type: none"> Work of merit, for example, some correct differentiation of $f(x)$ or $f'(x)$; states $f''(x) = 0$ Brings down derivative from (i) <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> Correct $f''(x)$ Finds x values of local maximum and local minimum

	$x = \frac{1}{2}$	
(c)	<p>Slope of $l = \frac{1}{2}$ Lines drawn parallel to l and touching the graph of $p(x)$ at $x \approx 2 \cdot 2$ and $x \approx 6 \cdot 8$</p>	<p>Scale 15D (0, 4, 6, 8, 15)</p> <p><i>Low Partial Credit</i></p> <ul style="list-style-type: none"> • Work of merit, for example, mentions slope of $l = \frac{1}{2}$ • Draws a line parallel to $l(x)$ • Draws a horizontal line at $y = \frac{1}{2}$ • Relevant work to draw graph of $p'(x)$ • Draws two parallel tangents to $p(x)$ that are not parallel to $l(x)$ <p><i>Mid Partial Credit</i></p> <ul style="list-style-type: none"> • One tangent drawn correctly <p><i>High Partial Credit</i></p> <ul style="list-style-type: none"> • Two tangents drawn correctly • One tangent drawn correctly and corresponding x value estimated correctly • Graphs of $l'(x)$ and $p'(x)$ shown on the diagram

Question 7

Q8	Model Solution – 50 Marks	Marking Notes
(a) (i)	<p>Top 10% means 90% below.</p> $P(z < 1.28) = 0.8997.$ $\frac{x-176}{36} = 1.28$ $\Rightarrow x = 222.08$ <p>Minimum mark of 223</p>	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p><i>Note:</i> Accept use of $P(z < 1.29)$, to give $x = 222.44$</p> <p><i>Note:</i> Accept answer rounded to 222 instead of 223</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Mean or standard deviation indicated • z-formula with some substitution <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • z-score found (1.28 or 1.29) • z-formula fully substituted $\left(\frac{x-176}{36}\right)$ <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • $\frac{x-176}{36} = 1.28$
(a) (ii)	$P(165 < x < 210)$ $P\left(\frac{165-176}{36} < z < \frac{210-176}{36}\right)$ $= P(-0.31 < z < 0.94) \text{ [2 D.P.]}$ $= P(z < 0.94) - P(z > -0.31)$ $P(z < 0.94) = 0.8264$ $P(z < -0.31) = 1 - P(z < 0.31)$ $= 1 - 0.6217 = 0.3783$ <p>So answer = $0.8264 - 0.3783 = 0.4481$</p> <p>= 44.81% of 1st years got the Distinction.</p>	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p><i>Note:</i> Also accept use of -0.30 instead of -0.31, and/or use of 0.95 instead of 0.94.</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Mean or standard deviation indicated • z formula with some substitution • $-0.305 \dots$ or $0.94 \dots$ <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • One relevant probability found directly from tables (0.6217 or 0.8264) <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • 0.3783 found • 0.8264 and 0.6217 found <p><i>Full Credit –1:</i></p> <ul style="list-style-type: none"> • Uses $P(z > -0.35)$ and finishes correctly

Q8	Model Solution – 50 Marks	Marking Notes
(b) (i)	$T = \frac{19.8 - 21}{\left(\frac{5.2}{\sqrt{60}}\right)} = -1.787 \dots$	<p>Scale 5C (0, 2, 3, 5)</p> <p>Note: Accept 1.787 ...</p> <p>Note: $\frac{s}{\sqrt{n}}$ must be used in order to be awarded above <i>Low Partial Credit</i></p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • Mean or standard deviation indicated • Relevant formula with some substitution <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • Formula fully substituted
(b) (ii)	<p><i>p</i>-value:</p> $p = 2[1 - P(z < 1.79)]$ $= 2(1 - 0.9633)$ $= 0.0734$ <p>Conclusion:</p> <p>There is not enough evidence to say that the claim in the news report is incorrect [as $0.0734 > 0.05$]</p>	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p>Note: Accept $P(z < 1.78)$, so <i>p</i>-value of $2(1 - 0.9625) = 0.075$.</p> <p>Note: Accept conclusion based on <i>z</i>-score rather than <i>p</i>-value.</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • $P(z < 1.79)$ • 0.9633 <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • $2[1 - P(z < 1.79)]$ • Work of merit in finding <i>p</i>-value and correct conclusion based on this <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • <i>p</i>-value found but no or incorrect conclusion
Q8	Model Solution – 50 Marks	Marking Notes
(c) (i)	<p>Assuming no replacement:</p> $\frac{18}{23} \times \frac{17}{22} \times \frac{16}{21} \times \frac{5}{20}$ $= 0.11518 \dots = 0.1152 \text{ [4 D.P.]}$ <p style="text-align: center;">OR</p> <p>Assuming replacement:</p> $\left(\frac{18}{23}\right)^3 \times \frac{5}{23}$ $= 0.10420 \dots = 0.1042 \text{ [4 D.P.]}$	<p>Scale 10D (0, 3, 5, 8, 10)</p> <p>Note: multiplication between relevant terms is necessary to be awarded above <i>Low Partial Credit</i></p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • One relevant fraction <p><i>Mid Partial Credit:</i></p> <ul style="list-style-type: none"> • Product of four fractions, two of them correct <p><i>High Partial Credit:</i></p> <ul style="list-style-type: none"> • $\frac{18}{23} \times \frac{17}{22} \times \frac{16}{21} \times \frac{5}{20}$ • $\left(\frac{18}{23}\right)^3 \times \frac{5}{23}$
(c) (ii)	$\left(\frac{12}{23} \times \frac{6}{22} \times \frac{5}{21}\right) 3! = \frac{360}{1771}$ $= 0.20327 \dots = 0.2033 \text{ [4 D.P.]}$ <p style="text-align: center;">OR</p> $\left(\frac{12}{23} \times \frac{6}{22} \times \frac{5}{21}\right) + \left(\frac{12}{23} \times \frac{5}{22} \times \frac{6}{21}\right) +$	<p>Scale 5C (0, 2, 3, 5)</p> <p><i>Low Partial Credit:</i></p> <ul style="list-style-type: none"> • One relevant fraction, for example, $\frac{12}{23}$ or $\frac{6}{22}$ or $\frac{5}{21}$ • Counts / lists different possible arrangements, for example, 3! or 6

$$\left(\frac{6}{23} \times \frac{12}{22} \times \frac{5}{21}\right) + \left(\frac{6}{23} \times \frac{5}{22} \times \frac{12}{21}\right) +$$

$$\left(\frac{5}{23} \times \frac{6}{22} \times \frac{12}{21}\right) + \left(\frac{5}{23} \times \frac{12}{22} \times \frac{6}{21}\right)$$

$$= 0.20327 \dots = 0.2033 \text{ [4 D.P.]}$$

OR

$$\frac{\binom{12}{1} \times \binom{6}{1} \times \binom{5}{1}}{\binom{23}{3}} = 0.2033 \text{ [4 D.P.]}$$

High Partial Credit:

- $\frac{12}{23} \times \frac{6}{22} \times \frac{5}{21}$ or any other relevant triple
- Assumes keys are replaced and finishes

Question 8

Q8	Model Solution – 50 Marks	Marking Notes
(a)	$F = 3000(1 + 0.024)^5$ $= \text{€}3377.70$	Scale 5B (0, 2, 5) <i>Partial Credit:</i> <ul style="list-style-type: none"> • Work of merit, for example, some correct substitution into relevant formula; finds 2.4% as a decimal
(b) (i), (ii)	<p>(i) It is the amount that should be invested today to amount to €1000 in 1 years' time at the particular interest rate.</p> <p>(ii) $4000 = P(1 + 0.024)^6$</p> $\frac{4000}{1.024^6} = P$ $P = \text{€}3469.45$	Scale 10D (0, 3, 5, 8, 10) Note: In (i) Accept $P = \frac{1000}{(1+i)}$ <i>Low Partial Credit:</i> <ul style="list-style-type: none"> • Work of merit in (i) or (ii), for example, formula in (i), correct substitution into relevant formula in (ii) • 2.4% written as a decimal <i>Mid Partial Credit:</i> <ul style="list-style-type: none"> • (i) or (ii) correct • Work of merit in both parts <i>High Partial Credit</i> <ul style="list-style-type: none"> • One part correct and work of merit in the other part
(c)	$1 \cdot 024 = (1 + i)^4$ $(1.024)^{\frac{1}{4}} = 1 + i$ $(1.024)^{\frac{1}{4}} - 1 = i$ $0 \cdot 005947.. = i$ Rate = 0.59%	Scale 5C (0, 2, 3, 5) <i>Low Partial Credit:</i> <ul style="list-style-type: none"> • Some correct substitution into relevant formula • 2.4% written as a decimal <i>High Partial Credit:</i> <ul style="list-style-type: none"> • $(1.024)^{\frac{1}{4}} = 1 + i$ • Evaluates correctly $i = (1.024)^4 - 1$ • Uses 3 or 12 instead of 4, but otherwise correct <i>Full Credit -1</i> <ul style="list-style-type: none"> • Answer given as a decimal
Q8	Model Solution – 50 Marks	Marking Notes
(d) (i) (ii)	<p>(i)</p> $A(1 \cdot 0011)^{36} + A(1 \cdot 0011)^{35} + \dots$ $\dots + A(1 \cdot 0011)^2 + A(1 \cdot 0011)$ <p style="text-align: center;">OR</p> $= A[(1 \cdot 0011)^{36} + (1 \cdot 0011)^{35} + \dots$ $\dots + (1 \cdot 0011)^2 + (1 \cdot 0011)]$ <p>(ii)</p> $A[1 \cdot 0011 + (1 \cdot 0011)^2 + \dots$ $\dots + (1 \cdot 0011)^{35} + (1 \cdot 0011)^{36}]$ $a = 1 \cdot 0011, \quad r = 1 \cdot 0011, \quad n = 36$ $A \left[\frac{1 \cdot 0011 (1 - 1 \cdot 0011^{36})}{1 - 1 \cdot 0011} \right] = 12000$ $A = \frac{12000}{1 \cdot 0011(1 - 1 \cdot 0011^{36})}$	Scale 15D (0, 4, 8, 12, 15) Consider as requiring 3 steps: 1. Finds geometric series 2. Substitutes into geometric formula 3. Finds A <i>Low Partial Credit:</i> <ul style="list-style-type: none"> • Work of merit in either part, for example, in (i) Writes 0.11% as a decimal; in (ii), sets answer in (i) equal to 12000 <i>Mid Partial Credit:</i> <ul style="list-style-type: none"> • 1 step correct • Substantial work of merit in both parts <i>High Partial Credit:</i> <ul style="list-style-type: none"> • 2 steps correct

	$1 - 1 \cdot 0011$ $A = €326 \cdot 60$ [2 D.P.]	<i>Full Credit –1:</i> <ul style="list-style-type: none"> • Correct solution, but excludes second and/or second last term • Investments made at the end of each month, otherwise correct
(e)	$E(x) = 11(0 \cdot 52) + (x - 5)(0 \cdot 15)$ $\quad \quad \quad + x(0 \cdot 33) = 13 \cdot 85$ $0 \cdot 15x + 0 \cdot 33x = 13 \cdot 85 - 5 \cdot 72 + 0 \cdot 75$ $0 \cdot 48x = 8 \cdot 88$ $x = €18 \cdot 50$	Scale 10C (0, 4, 7, 10) <i>Low Partial Credit:</i> <ul style="list-style-type: none"> • Work of merit, for example, some correct term in $E(x)$ <i>High Partial Credit</i> <ul style="list-style-type: none"> • Fully correct equation
(f)	<p>Cost Price = 82% of Selling Price Profit = 18% of Selling Price</p> $\text{Mark-up} = \frac{0 \cdot 18}{0 \cdot 82} \times 100$ $= 0 \cdot 2195 = 22\% \text{ [nearest percent]}$ <p style="text-align: center;">OR</p> <p>Let x = selling price and y = cost price</p> $\frac{x - y}{x} = 0 \cdot 18 \rightarrow y = 0 \cdot 82x$ <p>Mark up:</p> $\frac{x - y}{y} = \frac{x - 0 \cdot 82x}{0 \cdot 82x} = \frac{9}{41}$ <p>Mark up:</p> $\frac{9}{41} \times 100 = 21 \cdot 95$ $= 22\% \text{ [nearest percent]}$	Scale 5B (0, 2, 5) <i>Partial Credit:</i> <ul style="list-style-type: none"> • Work of merit, for example, states CP = 82% of SP • Mentions 82% • Finds 18% of a number